

University of Groningen

## The impact of galactic fountains on the global evolution of galaxy disks

Fraternali, F.; Binney, J.; Marasco, A.; Marinacci, F.

*Published in:*

Proceeding of the conference The Interplay between Local and Global Processes in Galaxies, Cozumel, Mexico, 2016-4, Eds. S. F. Sanchez, C. Morisset and G. Delgado-Inglada

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Publication date:*  
2016

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Fraternali, F., Binney, J., Marasco, A., & Marinacci, F. (2016). The impact of galactic fountains on the global evolution of galaxy disks. In *Proceeding of the conference The Interplay between Local and Global Processes in Galaxies, Cozumel, Mexico, 2016-4*, Eds. S. F. Sanchez, C. Morisset and G. Delgado-Inglada <http://adsabs.harvard.edu/abs/2016ilgp.confE..37F>

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*




# The impact of galactic fountain on disc evolution

Filippo Fraternali

Department of Physics and Astronomy, University of Bologna, I  
Kapteyn Astronomical Institute, University of Groningen, NL

L. Armillotta (Bologna), J. Binney (Oxford), A. Marasco (Groningen), F.  
Marinacci (MIT)

# Disc-corona interplay



Anderson & Bregman 2012

Dai+ 2012, Anderson+ 2013

MW Miller & Bregman+ 2013, 2015; Gatto+13

Mass corona  
~ 10-50% missing baryons



Cooling rate ~ 0.1 Mo/yr

# Global process: Fountain-driven gas accretion

# Disc-cloud corona interaction

$T_{\text{corona}} = 2 \times 10^6 \text{ K}$   $Z_{\text{corona}} = 0.1 Z_{\odot}$




$Z_{\text{cloud}} = 1 Z_{\odot}$



Mass of cold gas increased by  $\sim 20\%$ !



Marinacci, et al. 2010, 2011, MNRAS Lucia Armillotta

# Global fountain



Best-fit Accretion  
Rate  $\sim 3 M_{\odot} \text{yr}^{-1}$

Compare to SFR  $\sim 4 M_{\odot} \text{yr}^{-1}$



*Fraternali & Binney 2008, MNRAS; Marinacci, Fraternali+ 2011, MNRAS*



# Wav



0.5 0.7 0.9  
fraction


Compare to SFR  $\sim 1\text{-}3\text{ M}\odot\text{yr}^{-1}$

# Local process: Galactic hail: origin of the High Velocity Cloud complex C





# Origin of HVCs



**Oort 70** leftover of galaxy formation  
**Bregman 80** Galactic fountain  
 + satellites (**Olano 2001**), thermal instabilities  
 (**Kaufmann+ 06**), no thermal instabilities (**Binney+ 09**),  
 filaments (**Fernandez+ 12**)


**Wakker+ 1999, Nat**  $Z \sim 0.1$  Solar  $\rightarrow$  Accretion!

**Gibson+ 2001**  $Z \sim 0.3$  Accretion?

**Collins+ 2007** overabundance  $\alpha$  elements

(SN II?)

Filippo Fraternali (Bologna/Groningen)



Interplay local & global processes in galaxies - Cozumel, Mexico - 14/4/16

# Formation of complex C




Six free parameters:

$V_0$  = ejection velocity


$R_0$  = ejection location

$\delta R$  = size of the ejection




fraternali, Marasco, Armillotta, & Marinacci 2015, MNRAS Letter, 447, 70

# Origin & Location of complex



## Distance from the Sun




The distance is a completely independent confirmation

	$v_0$	$R_0$	$\delta_{\text{arm}}$	$t_0$	$\Delta t$	$t_{\text{dc}}$	(seed)	(final)
	[km s <sup>-1</sup> ]	[kpc]	[kpc]	[Myr]	[Myr]	[Myr]	[10 <sup>6</sup> M <sub>⊙</sub> ]	[10 <sup>5</sup> M <sub>⊙</sub> ]
Best-fit	211	9.5	2.9	150	53	46	3.4	6.8

$E_{\text{kin}} \sim 2 \times 10^{54}$  erg



# Hydrodynamical simulations



average metallicity at the end: **0.27 Solar**  
 compared to complex C: **0.1-0.3 Solar**

Complex C formed by a  
 superbubble + corona  
 condensation

*Local* manifestation of fountain-  
 driven accretion

## Trajectory of Smith Cloud

# Smith cloud



20 Million Years




Today

30 Million Years

NASA and ESA




Fox

Marasco, Fraternali+



# Conclusions

- Galactic fountain can cool the corona and feed the star formation in disc galaxies like the MW
- Local features like HVCs can form (condense) out of this non-linear perturbation of the corona
- Very good fit for the prototypical complex C, promising results for Smith cloud





# CROSSING THE RUBICON


The fate of gas flows in galaxies



<https://sites.google.com/site/rubiconf2016>

Santarcangelo di Romagna, Italy

5-9 September, 2016



## INVITED SPEAKERS



Manda Banerji (IoA, UK)  
Joel Bregman (Michigan Univ., USA)  
Natascha Förster-Schreiber (MPE, Germany)  
Joe Hennawi (MPIA, Germany)  
Andrew King (Leicester Univ., UK)  
Simon Lilly (ETH, CH)  
Federico Marinacci (MIT, USA)  
Raffaella Morganti (ASTRON, NL)  
Kate Rubin (CfA, USA)  
Jorge Sanchez-Almeida (IAC, Spain)  
Joop Schaye (Leiden Obser., NL)  
Francesco Tombesi (Goddard SFC, USA)  
Sylvain Veilleux (Maryland Univ., USA)  
Jessica Werk (UC S. Cruz, USA)

# Thanks




# Ophiucus superbubble

Purple HI  
Green Halpha



Energy to produce the holes:  
 $E \sim 1 \times 10^{53} - 1 \times 10^{55}$  erg




35m 20h34m10s  
Right Ascension (2000.0)


Boomsma et al. 2008,  
A&A

Pidopryhora et al. 2007,  
ApJ

# Our galactic fountain model




## Intermediate Velocity Clouds



Marasco, Fraternali & Binney 2012,

MNRAS

# Effect of spiral arms




Two limitations of our model:

1. *Axisimmetry*
2. *Average ejection velocities*


→ We introduced spiral arms



Disk gas



Fountain gas




Global  
model  
unchanged!


Best-fit  
Accretion  
Rate  $\sim 2$   
 $M_{\odot} \text{yr}^{-1}$



# NGC 6946



*Boomsma et al. 2007*





# SN-driven accretion in other sims

Modified SPH

No formation of clumps

*“Cold gas condenses from the halo at the intersection of supernovae-driven bubbles. This **positive feedback** feeds cold gas to the galactic disc*

*Habib et al. 2013, MNRAS*

MaGICC - GASOLINE

Halos enriched by galactic fountain

Gas in the fountain cycle comes back to the disk **more metal poor!**

*Brook+12, Brook+13*




# Impact of galactic fountain on disc evolution

1. Corona-disc interface
2. **Global process**: supernova-driven accretion
3. **Local process**: formation of condensed clouds
  - Origin of the high-velocity cloud complex C


# Positive feedback is there

$z=2$

Cooling induced close to galaxies by metals ejected by feedback



OWLS  
GADGET-3




*van de Voort & Schaye 2012*


# Ly $\alpha$ absorbers

HST/COS data


*Thom+ 2012, ApJL*



Impact parameters




Bound to the potential wells




# Early types vs star-forming

Cold gas ( $\log(T) < 5$ )



Thom+ 2012, *ApJL*

Hot gas ( $\log(T) \sim 5.5$ )



Tumlinson+ 2013

Werk+ 2013





Is this *cold* gas used  
for star formation?



# Fountain-driven accretion model



1. kick velocities ( $v_k$ )
2. Ionised fraction ( $f_{ion}$ )
3. Accretion rate



Fountain  
triggers coronal  
accretion at a  
rate  $\sim$  SFR

Best-fit Accretion Rate  $\sim 3 M_{\odot} \text{yr}^{-1}$


Compare to SFR  $\sim 4 M_{\odot} \text{yr}^{-1}$

*Fraternali & Binney,*

2008



# HI High Velocity Clouds



Typical  
Distances:  
 $\sim 10$  kpc

$h \sim \text{few-}10$  kpc

$Z \sim 0.1-0.4 Z_{\odot}$

$M < 10^7 M_{\odot}$

*Wakker et al. 2007, 2008; Tripp et al. 2003*

Accretion from High Velocity Clouds



$\sim 0.08 M_{\odot}/\text{yr}$



Includes He and factor 2 of ionised gas!

HI HVCs cannot feed  
SF

*Putman, Peek, Joungh 2012, ARA&A*

# Implications for galaxy evolution

# Global fountain




Best-fit Accretion Rate  $\sim 2 M_\odot/\text{yr}$

Compare to SFR  $\sim 1-3 M_\odot/\text{yr}$


Marasco, Fraternali+ 2012,  
MNRAS

Filippo Fraternali (Bologna/Groningen)




Interplay local & global processes in galaxies - Cozumel, Mexico - 14/4/16

# Metallicity



## Condensation of the corona




Supernova-driven  
cooling reproduces:  
Extraplanar HI  
Absorption features  
And even HVCs

Average metallicity at the end: **0.27 Solar**  
Compared to complex C: **0.1-0.3 Solar**



# Galactic fountain model

Building of several model cubes -> minimization residuals with LAB




We fit:

1. kick velocities ( $v_k$ )
2. Ionised fraction ( $f_{ion}$ )
3. Accretion coefficient ( $\alpha$ )


scaleheight  
vertical motions  
radial motions

$$\dot{m} = \alpha \dot{m}$$

## Pure fountain



## Fountain + accretion




# The effect of thermal conduction

Only cooling

$T_{\text{corona}} = 2 \times 10^6$  K


$Z_{\text{corona}} = 0.1 Z_{\odot}$

$Z_{\text{cloud}} = 1 Z_{\odot}$





Cooling & thermal conduction

$$\mathbf{F}_{\text{cond}} = f \times \kappa_{\text{Sp}} T^{5/2} \nabla T$$





# Galactic coronae





*Keres+ 2009*  
Hot-mode accretion. Similar to classical theory (e.g. *White & Rees 1978*)  
*& Bregman*

*2012*  
*Dai+ 2012, Anderson+ 2013*


MW *Miller & Bregman+ 2013, 2015; Gatto+13*  
Mass corona  
~ 10-50% missing baryons

Cooling rate ~ 0.1  $M_{\odot}/\text{yr}$




*Bogdan+ 2012*

# Disc-corona interplay



*Fraternali & Binney 2008, MNRAS*  
*Marinacci, et al. 2010, 2011, MNRAS*  
*Marasco, Fraternali & Binney 2012, MNRAS*


# Supernova-driven accretion



*Fraternali et al. 2013, ApJ*


# Efficiency of fountain-driven cooling

Evolution of cold gas mass



Convergence at  $\sim 2$  parsec


Evolution of cold gas mass



Armillotta, Fraternali & Marinacci, in prep



# Efficiency of fountain-driven cooling



Condensation strongly depends on coronal temperature

At  $T_{\text{cor}} > 4 \times 10^6 \text{ K}$  clouds evaporate

*Armillotta, Fraternali & Marinacci, in prep*

Condensation efficiency as a function of  $T_{\text{vir}} \leftrightarrow M_{\text{vir}}$

Role in the quenching of star formation?



$T_{\text{vir}}$

# Milky Way evidence

Chemical evolution models

G-dwarf problem

*Larson 1972, Tinsley 80,*



*Chiappini+ 97, 01; Schoenrich &*

*Binney 09*

Deuterium in local ISM appears  
to be re-supplied *Linsky et al.*

*2006*

Need for gas  
accretion at  $Z < \sim 0.1$




*Dauphas et al. 2005, Nature*

$$\text{SFR} \propto \dot{M}_{\text{acc}}$$

$$\sim 1 M_{\odot}/\text{yr}$$

*Fraternali & Tomassetti 2012, MNRAS*

# HI High Velocity Clouds



*Wakker et al. 2007, 2008; Tripp et al. 2003*

Typical  
Distances:  
 $\sim 10$  kpc

$h \sim \text{few-}10$  kpc

$Z \sim 0.1-0.4 Z_{\odot}$

$M < 10^7 M_{\odot}$

Accretion from High Velocity Clouds




$\sim 0.08 M_{\odot}/\text{yr}$

Includes He and factor 2 of ionised gas!

HI HVCs cannot feed  
SF

*Putman, Peek, Joungh 2012, ARA&A*

# Origin of HVCs



**Oort 70** leftover of galaxy formation


**Bregman 80** Galactic fountain

+ satellites (**Olano 2001**), thermal instabilities (**Kaufmann+06**), no thermal instabilities (**Binney+09**), filaments (**Fernandez+12**)

**Wakker+ 1999, Nature**  $Z \sim 0.1$  Solar  $\rightarrow$  Accretion!

**Gibson+01**  $Z \sim 0.3$  Accretion?

**Collins+07** overabundance  $\alpha$  elements (SN II?)






# Cosmology evidence

Gas depletion time  $\sim 1$  Gyr Assembly of stellar mass in the Universe


Gas depletion time  $t_{\text{depl}} = M_{\text{gas}} / \text{SFR}$



*Saintonge+ 15*

*Kennicutt+83, Genzel+ 10,*


*Bigiel+11, Genzel+15*



Compilation from *Madau & Dickinson 2014*


# Cosmology evidence

Constant HI in galaxies



Accretion needed to keep forming stars at level of  $\sim$  SFR

Bathtub model



*Zafar+ 13*

$$\text{SFR} \sim \dot{M}_{\text{acc}} - \dot{M}_{\text{outflow}} + \mathcal{R}\text{SFR}$$

*Bouché+2010, Davé+ 2012, Lilly+1*

# Supernova-driven accretion in other sims

# SN-driven accretion in other sims

Modified SPH

No formation of clumps

*“Cold gas condenses from the halo at the intersection of supernovae-driven bubbles. This **positive feedback** feeds cold gas to the galactic disc*

*Habib et al. 2013, MNRAS*

MaGICC - GASOLINE

Halos enriched by galactic fountain

Gas in the fountain cycle comes back to the disk **more metal poor!**

*Brook+12, Brook+13*







# Positive feedback is there

$z=2$

Cooling induced close to galaxies by metals ejected by feedback




OWLS  
GADGET-3





*van de Voort & Schaye 2012*

# Extrapolation in the Milky Way




Best-fit Accretion Rate  $\sim 2 M_{\odot}\text{yr}^{-1}$   
 Compare to SFR  $\sim 1\text{--}3 M_{\odot}\text{yr}^{-1}$



*Marasco, Fraternali & Binney 2012, MNRAS* Accretion in the outer disc

# Disc-corona interplay



Interface layer where  
disc and coronal  
material mix

↓  
**Cooling time** of the  
corona (typically very  
long) **decreases**  
**dramatically** because  
it is mixed with:

1. *cold* gas
2. High *Z* gas

*Fraternali & Binney 2008, MNRAS*

*Marinacci, et al. 2010, 2011, MNRAS*

*Marasco, Fraternali & Binney 2012, MNRAS*

# Implications for galaxy evolution





# Evolution of discs

Red and dead




And the corona does not cool further

If gaseous disc is too hot







Cold mode






Dekel et al. 2009

# Efficiency of SN-driven cooling



# Conclusions



- Supernova feedback cools the corona in star-forming galaxies like the MW
- **Very good fits:**
  - HI in the MW and external galaxies,
  - ionized absorbers in the MW
- This can be the way hot-mode accretion feeds star formation in disc galaxies




# 1. Extraplanar HI in the Milky Way



# HI disk and halo in the Milky Way



# HI halo – all-sky








*Marasco & Fraternali 2011, A&A*

Pure fountain

Fountain + accretion

HI data

Best fit



$v_k = 75 \text{ km/s}$   
 $f_{\text{ion}} = 0.3$

$\dot{M}_{\text{cor}} \sim 2 M_{\odot}/\text{yr}$



Halo gas:  
 $\sim 80\%$  from fountain  
 $\sim 20\%$  from corona



# 2. Absorption features



# Cooling in the wake




*Fraternali et al. 2013, ApJL*

*Shull+ 2009, ApJ*



*Lehner & Howk 2011, Science*

*Lehner et al. 2012, MNRAS*

C II, Si II, Si III, ...  $4.3 < \log T < 5.3$   
K



# Evolution of the wake



DYNAMICAL MODEL  
for the IONISED GAS  
(no free parameters)




Marasco, Marinacci & Fraternali 2013, MNRAS

Filippo Fraternali (Bologna/Groningen)

Interplay local & global processes in galaxies – Cozumel,  
Mexico – 14/4/16

# in the MW

larasco



Data from Lehner et al. 2012, MNRAS

This model reproduces:

- Positions & velocities of **95% absorbers**
- Average column density
- Number of absorbers along the l.o.s.
- **High velocity dispersions** of absorbers

‘Warm’ accretion:  $\sim 1 M_{\odot}/\text{yr}$

Filippo